

Radial Velocity Studies of Close Binary Stars. X¹

Slavek M. Rucinski

*David Dunlap Observatory, University of Toronto
P.O. Box 360, Richmond Hill, Ontario, Canada L4C 4Y6*

`rucinski@astro.utoronto.ca`

Wojtek Pych

*David Dunlap Observatory, University of Toronto
and Copernicus Astronomical Center, Bartycka 18, 00-716 Warszawa, Poland*

`pych@camk.edu.pl`

Waldemar Ogłóza

*Mt. Suhora Observatory of the Pedagogical University
ul. Podchorążych 2, 30-084 Cracow, Poland*

`ogloza@ap.krakow.pl`

Heide DeBond, J. R. Thomson, Stefan W. Mochnacki, Christopher C. Capobianco²,
George Conidis

*David Dunlap Observatory, University of Toronto
P.O. Box 360, Richmond Hill, Ontario, Canada L4C 4Y6*

`(debond,jthomson,mochnacki,conidis)@astro.utoronto.ca, ccapo@ap.smu.ca`

P. Rogoziecki

Adam Mickiewicz University Observatory, Słoneczna 36, 60-286 Poznań, Poland

`progoz@moon.astro.amu.edu.pl`

ABSTRACT

Radial-velocity measurements and sine-curve fits to the orbital velocity variations are presented for the ninth set of ten close binary systems: V395 And, HS Aqr, V449 Aur, FP Boo, SW Lac, KS Peg, IW Per, V592 Per, TU UMi, FO Vir. The first three are very

²Current address: Department of Astronomy and Physics, Saint Mary's University, Halifax, Nova Scotia, Canada B3H 3C3

close, possibly detached, early-type binaries and all three require further investigation. Particularly interesting is V395 And whose spectral type is as early as B7/8 for a 0.685 day orbit binary. KS Peg and IW Per are single-line binaries, with the former probably hosting a very low mass star. We have detected a low-mass secondary in an important semi-detached system FO Vir at $q = 0.125 \pm 0.005$. The contact binary FP Boo is also a very small mass-ratio system, $q = 0.106 \pm 0.005$. The other contact binaries in this group are V592 Per, TU UMi and the well known SW Lac. V592 Per and TU UMi have bright tertiary companions; for these binaries, and for V395 And, we used a novel technique of the broadening functions arranged into a 2-dimensional image in phase. The case of TU UMi turned out intractable even using this approach and we have not been able to derive a firm radial velocity orbit for this binary. Three systems of this group were observed spectroscopically before: HS Aqr, SW Lac, KS Peg.

Subject headings: stars: close binaries - stars: eclipsing binaries – stars: variable stars

1. INTRODUCTION

This paper is a continuation in a series of papers (Papers I – VI and VIII – IX) of radial-velocity studies of close binary stars (Lu & Rucinski 1999; Rucinski & Lu 1999; Rucinski, Lu, & Mochnacki 2000; Lu, Rucinski, & Ogloza 2001; Rucinski et al. 2001, 2002, 2003; Pych et al. 2004) and presents data for the ninth group of ten close binary stars observed at the David Dunlap Observatory. For technical details and conventions, and for preliminary estimates of uncertainties, see the interim summary paper Rucinski (2002a, Paper VII). Selection of the targets is quasi-random: At a given time, we observe a few dozen close binary systems with periods shorter than one day, brighter than 10 – 11 magnitude and with declinations $> -20^\circ$; we publish the results in groups of ten systems as soon as reasonable orbital elements are obtained from measurements evenly distributed in orbital phases.

Whenever possible, we estimate spectral types of the program stars using our classification spectra. These are compared with the mean $(B - V)$ color indexes taken from the *Tycho-2* catalog (Høg et al. 2000) and the photometric estimates of the spectral types using the relations published by Bessell (1979).

The observations reported in this paper have been collected between June 1997 and November 2004. The ranges of dates for individual systems can be found in Table 1. All systems discussed in this paper, except three: HS Aqr, SW Lac, and KS Peg, have been observed by us for radial-velocity variations for the first time. We have derived the radial velocities in the same way as described in previous papers. See Paper VII for a discussion of the broadening-function approach used in the

¹Based on the data obtained at the David Dunlap Observatory, University of Toronto.

derivation of the radial-velocity orbit parameters: the amplitudes, K_i , the center-of-mass velocity, V_0 , and the time-of-primary-eclipse epoch, T_0 .

The novelty of this paper is the treatment of binaries with very weak signatures of binarity in their broadening functions, resulting either from exceptionally weak Mg I $\lambda 5184$ triplet lines (V395 And, Section 2.1) or from the presence of a bright third star in the system (TU UMi, Section 2.9) which limits the dynamic range in the broadening function. We utilize here an approach borrowed from image processing techniques in that we represent the broadening function information in the phase domain as a two-dimensional image where smoothing in the phase coordinate restores the temporal correlation between successive broadening functions.

This paper is structured in a way similar to that of previous papers, in that most of the data for the observed binaries are in two tables consisting of the radial-velocity measurements in Table 1 and their preliminary sine-curve solutions in Table 2. Table 1 contains data for nine stars as we have not been able to derive individual radial velocities for the difficult case of TU UMi.

The data in Table 2 are organized in the same manner as in previous papers. In addition to the parameters of spectroscopic orbits, the table provides information about the relation between the spectroscopically observed epoch of the primary-eclipse T_0 and the recent photometric determinations in the form of the $O - C$ deviations for the number of elapsed periods E . It also contains our new spectral classifications of the program objects. Section 2 of the paper contains brief summaries of previous studies for individual systems and comments on the new data. Figure 1 and Figure 4 show the 2-D broadening function images for V395 And and TU UMi while Figure 2 and Figure 3 show the radial velocity data and solutions. Figure 5 shows the BF's for all systems; the functions have been selected from among the best defined ones, usually around the orbital phase of 0.25 using the photometric system of phases counted from the deeper eclipse.

2. RESULTS FOR INDIVIDUAL SYSTEMS

2.1. V395 And

Photometric variability of the star was discovered by the Hipparcos satellite (ESA 1997); the star was assigned RRc type in the 74th Special Name-list of Variable Stars (Kazarovets et al. 1999) on the basis of the Hipparcos data. Duerbeck (1997) suggested that V395 And is a contact eclipsing system on the basis of the period-color relation. Light elements were calculated by Pribulla et al. (2003), HJD=2448500.4372 +E 0.684656 using the Hipparcos photometric data and the doubled value of the original period.

V395 And was listed in the Duflot et al. (1995) with the spectral type of this object estimated as B8 (this type is also quoted in the Washington Double Star Catalog², from now on called WDS).

²The Washington Double Star Catalog is available at the Internet site: <http://ad.usno.navy.mil/wds/>.

and radial velocity of -8 km s^{-1} ; in the HD catalog the spectral type is A0. Our spectra of the magnesium triplet in V395 And showed an exceptionally weak, poorly defined, strongly broadened feature with depth of only 2 percent of continuum with no discernible doubling. While we expected that the feature will be weak in a nominally A0 type star (Simbad), the particular weakness of the line was surprising.

The star is relatively bright, $V_{max} = 7.55$. It is a member of a visual binary (WDS 23445+4623). The companion at a separation of 5.7 arcsec is 2.2 mag fainter than V395 And. It is visible at the position angle 104° and thus is projected into our East–West spectrograph slit. Its presence may have produced weakening of the spectral lines by probably less than 15%.

Original estimates based on the very shallow and poorly defined broadening functions (BF’s) suggested that the star will be entirely intractable and may have to be abandoned. Our standard technique of measuring radial velocities requires that each BF, for each spectrum, gives a peak or a pair of peaks for radial velocity centroid determination; such well defined peaks simply did not exist for V395 And. However, a new approach, partly based on ideas used in image processing worked relatively well: The BF’s were ordered in as columns in a two-dimensional image, with Y-coordinate being the heliocentric radial velocity. The X-coordinate, the phase, is obviously unevenly filled and the data can be interpolated into a regular grid of phases. At this point, we utilize the fact that the BF’s are correlated in the temporal (phase) domain so we can use smoothing. Thus, horizontal cuts through the BF-image were convolved with a Gaussian kernel with the full width at half maximum (FWHM) of 0.025 in phase, and then re-binned into a new uniform phase system with 40 phase points. The “broadening function image” is shown in Figure 1 while a section at phase 0.75 is shown in Figure 5, among the broadening functions for other stars of this group.

Even after processing of all available 103 spectra of V395 And, as described above, only some of the smoothed BF’s were measurable for individual radial velocities. They are listed in Table 1 as for other stars, but they should be used with considerable caution. The smoothed broadening functions were originally found at 0.025 intervals in phase and heliocentric times were calculated for them using the T_0 prediction given above. Some of the smoothed BF’s were still too poor and have been eliminated from the listing. The final solution is based on the data as listed in Table 1 and required a new value of the initial epoch T_0 , as given in Table 2.

The star appears to be a very close pair with almost identical components, $q = 0.88 \pm 0.03$. The component potentially eclipsed at the primary minimum (corresponding to T_0) is less massive; its peak in the broadening function is slightly sharper and taller indicating higher surface temperature and lower rotation rate. The very small amplitude of light variations, $\Delta V \simeq 0.04$, suggests that the eclipses will not be actually observed and that the binary is an “ellipsoidal variable” showing only proximity effects. Our classification spectra clearly indicated a spectral type as early as B7 which is difficult to reconcile with the close orbit with the period of 0.685 days. The two spectra that we obtained were not identical and showed some peculiar variability and emission at He I $\lambda 4009/4027$. No spectral estimate of $\log g$ was possible due to peculiarity of the spectra. Stars of the spectral

type of B7 or B8 have masses around 2.8 to 3.0 $M_{\odot}$ and radii around 2.2 to 2.4 $R_{\odot}$. If the orbit has $i \simeq 40^{\circ}$ then the current data would imply $A \simeq 6.0 R_{\odot}$ which is just about possible for a contact system.

2.2. HS Aqr

HS Aqr is a detached or semi-detached binary. Spectroscopic studies based on the high quality echelle spectra were presented by Popper (1996, 1997, 2000). This was followed by an extensive discussion of the photometric and spectroscopic data by Clausen et al. (2001). The above sources provide the most extensive references to properties of the system. The value of T_0 was taken from Kreiner (2004): $T_0 = 2,452,500.6960 + E \times 0.710188$.

Our observations of HS Aqr give a good solution of the radial velocity orbit (Figure 2), but perhaps not of such a high quality as that of Popper which was based on the echelle spectrograph spectra. However, since the main uncertainties in this field are not in random errors, but in systematic differences in measuring techniques, we give our solution here for future reference.

The broadening function at phase close to 0.25 (Figure 5) shows two well defined peaks for detached components, with that for the secondary components a bit broader indicating that this component is larger, if this star is in rotation-revolution synchronism. This agrees with the supposition (Clausen et al. 2001) that the secondary component is close to or fills its Roche lobe.

Our new spectral type estimate of F6V is slightly earlier than previously discussed (F8V and G8-9), in a better agreement with the Tycho-2 (Høg et al. 2000) color index $B - V = 0.52$. The binary is in a visual system (CCDM J20409-0036AB, ADS 14147AB), but the companion at 1.6 arcsec separation is some 4 magnitudes fainter than HS Aqr thus of no consequence for the radial velocity orbit. The recent minimum prediction by Kreiner (2004) perfectly agrees with our determination of the initial epoch T_0 .

The star is bright, $V_{max} = 9.0$, but was not included in the Hipparcos database, so its parallax is currently unknown.

2.3. V449 Aur

V449 Aur was discovered photometrically by the Hipparcos mission. The primary eclipse prediction was the HIP one: $T_0 = 2,448,500.2670 + E \times 0.703648$. The mean *uvby* data of Jordi et al. (1996), $V = 7.455$ and $b - y = 0.086$, suggest an early-type no later than A5/6V, while our spectral classification is A2IV and, through a comparison with the color index, implies some reddening.

The binary type of the system is not clear at this moment: The broadening functions are par-

tially blended, as in a contact binary, and somewhat difficult to measure for centroid determinations (see Figure 5). On the basis of its very shallow Hipparcos light curve, but with rather well-defined eclipses, the binary was considered by Rucinski (2002b) to be a detached one (EA); the broadening functions would rather suggest a contact binary. However, the large widths of the peaks in the broadening functions and the resulting blending may result from rotational velocities higher than the synchronous ones. The mass ratio is not far from unity, $q = 0.730 \pm 0.017$, which is frequently encountered among early-type very close binaries. The scatter of the radial velocity observations of the slightly more massive component, which is eclipsed in the primary minimum, appears to be genuinely larger at phases around 0.2 – 0.4 (Figure 2); we have no explanation for it.

2.4. FP Boo

FP Boo is another case of the Hipparcos photometric discovery. The Hipparcos period and T_0 serve the current observations relatively well: $T_0 = 2,448,500.478 + E \times 0.640487$.

The previous spectral type given by SIMBAD and in the Hipparcos database, A5, does not agree with our estimate of F0V; the latter would imply $B - V = 0.28$ whereas the Tycho-2 mean color index is even redder, $B - V = 0.35$, corresponding to F2V.

The Hipparcos photometry was analyzed by Selam (2004) using a simplified approach. The mass ratio was estimated at $q_{ph} = 0.1$, which is in a surprisingly good agreement with the spectroscopic result, $q_{sp} = 0.106 \pm 0.005$, taking into account that the light curve is shallow and does not show any obvious total eclipses. (This case is – in our view – an exception, a fortuitous coincidence. Normally, we do not trust the so-called photometric mass ratios because so many stars in our RV program show $q(sp) \neq q(ph)$). FP Boo was somewhat faint ($V_{max} = 10.07$) for Hipparcos photometry, so the light curve could be easily improved. Since the velocity amplitudes are large and thus imply the orbital inclination close to 90 degrees, it is likely that more precise photometry will show total eclipses. In its small mass ratio and the spectral type of F0/2V, the system is somewhat similar to the well-known contact binary AW UMa.

The radial velocity orbit is shown in Figure 2 while a broadening function close to phase 0.25 is shown in Figure 5.

2.5. SW Lac

SW Lac is one of the most frequently photometrically observed contact binaries. It is a bright ($V_{max} = 8.66$) binary with a large amplitude and an unstable light curve and thus an easy target for investigations utilizing small telescopes. Several discussions addressed the matter of evolving surface spots, the most recent being by Albayrak et al. (2004) where references to previous work can be found.

SW Lac is a late type contact binary with the spectral type variously assigned as G3, G5 and K0. Our new classification is G5V which predicts a color index a bit less red than the observed $B - V = 0.73$; possibly G8V would be a good compromise. From the moment of minima predictions as well as the direct detection of faint spectral signatures (Hendry & Mochnacki 1998), the binary appears to have at least one – probably two – nearby, low mass companions (Pribulla et al. 1999). The system is known to change the orbital period. We used the moment of primary minimum of Kreiner (2004): $HJD = 2,452,500.1435 + E \times 0.3207158$ which ideally predicts our T_0 .

Our observations form the third currently available set of radial velocity data (see Figure 2). Zhai & Lu (1989) published the first radial velocity orbit, with $q = 0.797 \pm 0.010$. A similar, but not identical result was obtained soon after by Hrivnak (1992) who published only the mass ratio, $q = 0.73 \pm 0.01$, without more details. In spite of the large brightness of the star and excellent definition of the individual peaks in the broadening functions, our mass ratio, $q = 0.776 \pm 0.012$, carries a larger error than for most of our targets, probably because of the genuine changes in the broadening functions, which are caused by elevated stellar surface activity. What is important, however, that radial velocity semi-amplitudes in our orbit (Figure 2) are larger than previously observed: Our total mass estimate, $(M_1 + M_2) \sin^3 i = 2.10 \pm 0.06 M_\odot$ (the primary $2.18 M_\odot$, the secondary $0.92 M_\odot$), is significantly larger than for the orbit of Zhai & Lu (1989), $(M_1 + M_2) \sin^3 i = 1.74 M_\odot$. Since most of systematic effects result in a reduction of the radial velocity amplitudes, our result is more trustworthy as it is based on the superior technique of the broadening functions. Thus, SW Lac is a relatively massive, but surprisingly cool system, a situation which was very clearly seen before in the case of another cool contact binary AH Vir (Lu & Rucinski 1993). As in the case of AH Vir, the masses are characteristic for F-type stars while the combined luminosity is in agreement with small star sizes – implied by the Roche geometry of a very short period system – and their low temperature (G5V). Indeed, for SW Lac, the M_V calibration (Rucinski & Duerbeck 1997) predicts $M_V(cal) \simeq 4.5$ as compared with the parallax-based, observed $M_V(obs) = 4.1 \pm 0.2$, so that the geometry and photometric properties are those of a late-type, compact contact system, only the masses are too large.

We have not noticed in individual broadening functions any traces of the third component detected by Hendry & Mochnacki (1998) and apparently producing part of the time-of-minima perturbations (Pribulla et al. 1999; Kreiner 2004). However, the same spectra, when averaged in the heliocentric system, clearly show a late type dwarf contributing about 1 percent to the total light at our 5184 Å spectral window (Caroline D’Angelo and Marten van Kerkwijk, 2005, private communication).

2.6. KS Peg

The bright star KS Peg ($V_{max} = 5.48$, $B - V = +0.01$), also called 75 Peg, has been known as a radial velocity variable for a long time, since the work of Plaskett et al. (1920). Hube & Gulliver (1985) recognized the peculiar properties of this binary star, in particular the very small mass

ratio of $q < 0.1$ making the velocity amplitude of the primary component small and the secondary component invisible in their spectra. The photometric variability was observed by Hube et al. (1988) while a good quality light curve was provided by the Hipparcos (ESA 1997) project. The light curve with amplitude of only 0.1 magnitude shows very wide minima with unusual, sharply peaked, brief light maxima.

We observed KS Peg in our standard way, and – because the star is so bright – with a particular attention to detection of the secondary component. In our series of observations, we have already detected several binaries with $q \simeq 0.1$, with some even below this limit, such as SX Crv at $q \simeq 0.07$ (Rucinski et al. 2001), so we hoped to see a signature of the secondary. The spectral type of KS Peg is A1V; we used the star of A2V as a main template, but we also used another F8V radial velocity standard to possibly capture the signature of the secondary component of later spectral type. None of these led to detection of any traces of the secondary, even when applying the technique of the “BF-image” smoothing described above for V395 And (Section 2.1) which permits detection of weak features correlating in the phase domain. Since the spectra of such a bright star result in particularly good determinations of the broadening functions, the non-detection of the secondary is puzzling and indicates that the secondary may be a very low mass star or even a massive planet, just able to enforce rapid rotation on the visible component, but too faint to contribute light to the overall luminosity of the binary.

One of the panels of Figure 5 shows comparison of broadening functions at both orbital elongations (averages of 6 BF’s at each elongation). Rapid rotation of the primary component and its (small) orbital displacements are very clearly visible, but there is no sign of the secondary component. Note that systematic trends in the baselines are due to the rectification process with inevitably different tracing of the continuum for the program and the standard template spectra. These trends may mask a very weak secondary component.

Our initial reference epoch was that of the Hipparcos photometry, $T_0 = 2,448,500.286 + E \times 0.502103$, which predicts our T_0 very well. In fact, with the same period as in the table, even the epoch of the Hube & Gulliver (1985) is well predicted more than 16 thousand epochs earlier (the authors in fact suggested 0.5021035, which does not tie all observations that well). The remarkable constancy of the orbital period may be one of useful hints in interpretation of KS Peg.

KS Peg is an important object because it is a bright star and thus belongs to the complete, magnitude-limited Hipparcos sample of short-period binaries brighter than 7.5 magnitude (Rucinski 2002b). Its parallax is moderately large, 13.65 ± 0.75 milli-arcsec, and the star is intrinsically bright, $M_V = +1.1 \pm 0.1$. The absolute magnitude agrees very well with the expected value for a Main Sequence A1 star, $M_V \simeq +1.0$. The large projected rotational velocity of the primary component, $V_1 \sin i = 240 \pm 15 \text{ km s}^{-1}$, implies that the orbital inclination cannot be far from 90° degrees. Assuming $M_1 = 2.3 \pm 0.1 R_\odot$, $R_1 = 2.3 \pm 0.1 M_\odot$, one can achieve a consistent picture, in terms of reproduction of K_1 and $V_1 \sin i$, with a relatively small secondary star with $M_2 = 0.18 \pm 0.01 M_\odot$ and thus $q = 0.078 \pm 0.005$, in a binary with the orbital inclination $i = 80^\circ \pm 3^\circ$.

2.7. IW Per

IW Per is somewhat similar to KS Peg described above: It is also bright, $V_{max} = 5.76$, small amplitude, 0.04 mag., photometric variable. The variability was recognized by Morris (1985), but most of the research concentrated on the chemical peculiarities and Am characteristics (Abt & Morrell 1995; Adelman 1998; Pauzen & Maitzen 1998).

IW Per was not a lucky star in our observations. A new CCD system which was used for it failed after some time and the data could not be related to those taken before or after as our technique requires high consistency in observations of program and template stars. Later on, it was realized, that the heavy dewar of the CCD system induced flexure in the spectrograph of up to 1/2 of the pixel in the detector focal plane. We decided to analyze and publish the available data for IW Per when it was realized that it is a single-line spectroscopic binary, precluding any in-depth studies of this system. In the final radial velocity solution, the spectra taken at large hour angles have been given half weight to account for the spectrograph flexure. The period and the initial epoch for our observations have been taken from the Hipparcos database. $T_0 = 2,448,500.523 + E \times 0.917180$.

In contrast to KS Peg, IW Per is not a very low amplitude radial velocity variable: The half amplitude $K = 98.6 \text{ km s}^{-1}$ indicates that the mass ratio is not very small, probably around $q \simeq 0.45 - 0.5$ and that the spectroscopically invisible component is a star with a mass only slightly smaller than solar. The visible component rotates with $V_1 \sin i \simeq 90 \pm 10 \text{ km s}^{-1}$ which is consistent with synchronous rotation in a moderately wide binary with the orbital period of 0.917 days.

IW Per has the best determined parallax among stars of the current group, $18.29 \pm 0.81 \text{ mas}$. The implied luminosity, $M_V = 2.07 \pm 0.11$, the spectral type A3V and $B - V = 0.13$ are mutually consistent.

2.8. V592 Per

V592 Per is another photometric discovery of Hipparcos. It is a relatively bright, $V_{max} = 8.22$, basically unstudied contact binary. The moment of primary eclipse from Hipparcos, $HJD = 2,448,500.243 + E \times 0.715722$ serves well for prediction of T_0 . The star has been known to possess a visual companion at a separation of only 0.16 arcsec (Heintz 1990). Data for WDS 04445+3953 indicate that the position angle changes rather rapidly so the system is definitely a physical one. We re-discovered the companion spectroscopically: In the broadening functions, its signature is strong, because of its sharpness, but the companion is fainter than the binary with $L_3/(L_1 + L_2) = 0.60 \pm 0.06$, or $\Delta m = 0.55 \text{ mag}$; in the WDS catalog $\Delta m = 0.85 \text{ mag}$.

To determine the close binary orbit, we analyzed our observations the way as before for triple systems, by first fitting three Gaussians to the whole BF, then subtracting the third star Gaussian and then measuring the radial velocities of both binary components from the residual broadening functions. In spite of the presence of the visual companion, the contact binary orbit is well defined,

with $q = 0.408 \pm 0.008$. The spectroscopic result is in total disagreement with the mass ratio estimated from the light curve of $q_{ph} = 0.25$ (Selam 2004). Once again, we stress that – due to the very low information content of light curves – light curve solutions for partially eclipsing contact binaries have little sense and they should not be attempted without supporting spectroscopic data.

Our observations are spread over time in that there is a gap of two full seasons. The radial velocities of the third component were estimated from all available spectra giving the average value of $V_3 = +29.28 \pm 0.20$ km s⁻¹ for the first season (2,451,870 – 2,451,957) and $V_3 = 27.92 \pm 0.18$ km s⁻¹ for the second season (2,452,513 – 2,452,517). Our systematic uncertainties are at the level of 1.5 – 2.0 km s⁻¹, while measurements in the presence of the variable binary “background” difficult, so we cannot claim that we see seasonal variation in V_3 ; it is basically identical to V_0 of the contact binary.

In view of similar luminosities, the spectral type F2IV estimated by Grenier et al. (1999) most probably applies to both stars, the contact binary V592 Per and the third star in the system. This spectral type does not agree with $(B - V) = 0.47$ estimated from the Tycho-2 data (Høg et al. 2000) which suggests F5–F6.

2.9. TU UMi

TU UMi is one of the Hipparcos photometric discoveries. A description of new photometric observations conducted concurrently with the spectroscopic observations was given by Pych & Rucinski (2004); the paper contains also references to work related to identification of the variability and of the determination of the period at twice the Hipparcos original period.

The previous photometric paper (Pych & Rucinski 2004) was based on a short time interval and could not provide a new value of the period which was estimated to only 4 significant figures. However, the Hipparcos period, which we use here, together with the new photometric epoch, $JD(\text{min}) = 2,452,725,6262$ (Pych & Rucinski 2004), counted back to the Hipparcos epoch (additionally shifted by 1/4 of the period because the star was thought to be a pulsating one so the time of its maximum light was used), $JD(\text{min}) = 2,448,499.9747$, indicate a good stability of the period.

TU UMi is a case of a contact binary with a bright close companion. We were not aware that it had been known as a visual binary WDS 14557+7618 with separation of 0.2 arcsec (this is well below the 1.88m telescope resolution in our location) when we started the observations. The spectral lines of the close binary are very broad and, in fact, entirely invisible in individual spectra, being masked by the strong sharp spectrum of the companion.

With 70 individual spectra rather evenly covering orbital phases, we attempted to analyze the data using the approach similar to that described for V395 And (Section 2.1) by ordering in phase and smoothing the broadening functions. This case is much more difficult than that of V395 And

because, even after averaging, the binary features are too weak and too poorly defined for being measured; most of the dynamic range in the BF is “wasted” on the sharp peak of the tertiary component (Figure 5). We attempted to remove this peak from the broadening function. First, we followed our normal routines of approximating the third star feature by a Gaussian and subtracting it. This did not work as the profile is not Gaussian at its base where its shape is particularly important. Then we attempted to find the shape of the third-star peak by combining all broadening functions and determining their lower envelope. This still did not provide a correct shape of the tertiary peak at the base, since none of the BF’s was really free of the binary signature. Thus, we were unable to remove the tertiary and to determine individual radial velocities of components for TU UMi; this star does not have any data listed in Table 1. While this is disappointing, we feel that it is doubtful if this star will ever permit a detailed parameter determination. In this situation, we present only very rough estimates from what can be derived from just eyeballed fit to the broadening functions in the 2-dimensional representation (Figure 4).

In plotting the expected changes in the broadening functions as a function of phase, we assumed the initial epoch as observed simultaneously photometrically by Pych & Rucinski (2004). The star appears to be a W-type contact system with a rather large amplitude of the secondary star variation, perhaps as large as $K_2 \simeq 220 \pm 20 \text{ km s}^{-1}$. An estimate for the primary is $K_1 \simeq 35 \pm 15 \text{ km s}^{-1}$. These amplitudes are shown on the BF image in Figure 4. Integrations of the wide binary feature at several phases indicate that the binary is in fact only marginally fainter than the third component, $L_3/(L_1 + L_2) = 1.25 \pm 0.15$, but the sharpness of its peak dominates in the combined spectrum of the system. The above estimate of the relative luminosity may carry a large systematic error since it is strongly dependent on how the baseline of the broadening function is drawn; in fact, this luminosity ratio implies $\Delta m = -0.24 \pm 0.12$, which poorly agrees with the direct magnitude difference given in the WDS catalog ($\Delta m = -0.47$).

The third component was observed to have the mean velocity of $-4.16 \pm 0.20 \text{ km s}^{-1}$ (a single observation $\sigma = 1.7 \text{ km s}^{-1}$).

2.10. FO Vir

FO Vir is a totally eclipsing, detached or semi-detached binary, with the larger component close to filling the Roche lobe, thus possibly an object in the pre-contact state. An extensive discussion of its properties, following the photometric study of Schmidt & Fernie (1984) who established its binary character, was presented by Mochnacki et al. (1986). On the basis of photometry and radial velocities of the dominant A7V component, they were able to infer many parameters of the system, in particular they predicted the mass ratio, $q = 0.15 \pm 0.02$, in a semi-detached system with the more massive component filling or almost filling its Roche lobe. Similar results were obtained by Poretti et al. (1987).

We have been able to detect the secondary component of FO Vir, but with some difficulty,

because it is relatively small and faint at the observed mass ratio $q = 0.125 \pm 0.005$ and much later spectral type than that of the A7V primary, probably early K. A somewhat indirect detection of the secondary was reported by Shaw et al. (1991) who observed the Lyman-alpha $\lambda 1216$ emission moving in anti-phase with the primary star and thus most likely related to the secondary; their $q = 0.16 \pm 0.04$ is in agreement with the current determination.

We experienced difficulties with measuring the radial velocities of both components in the first half of the orbit; this could be related to the previously reported strong and variable O’Connell effect (Shaw et al. 1991). In addition, the case of FO Vir confronted us with what may be a limitation of our broadening functions approach: We observed relatively strong (similar to those of the secondary component), initially unexplained broadening-function features moving in phase with the primary. We suspect that these “ghosts” may appear because of the low metallicity of the primary component (Mochnecki et al. 1986) and a mismatch between the spectra of program and template stars; we simply could not locate a proper, low metallicity template among bright standard stars. Many tests and experiments with different templates give us assurance that the final orbital elements are secure ones, although certain element of uncertainty still remains.

The radial velocity data for FO Vir may have to be re-interpreted when a better T_0 prediction for the duration of our observations (1997 – 2001) becomes available. With observations spanning four seasons, we could establish that the period must be shorter than that used by the Hipparcos project (0.775569) and we used one which is only by one digit in the last place shorter than that; however, the period may be – in fact – systematically changing so that a full discussion is clearly in order, especially in view of a possibly sporadic mass transfer in the system. Our T_0 is shifted by 50 minutes from the time predicted by the Hipparcos project. A re-discussion of rather few available moments of minima for 1983 – 1991 gave $JD(hel) = 2,452,500.1514 + E \times 0.7755678$, which we took as a guidance for adopting a shorter period, but which nevertheless resulted in the spectroscopic T_0 for our program too early by 40 minutes.

FO Vir is a member of a visual binary WDS 13298+0106, but the companion is some 6.5 fainter so that its presence was of no consequence for the spectroscopic observations.

3. SUMMARY

This paper presents radial-velocity data and orbital solutions for the ninth group of ten close binary systems observed at the David Dunlap Observatory. This group contains very interesting objects: KS Peg and IW Per are single-line (SB1) binaries without any traces of companions; the former may have a very small-mass secondary ($q < 0.1$). V395 And, HS Aqr and V449 Aur are very close, early-type binaries; the matter of them being detached systems requires further work; V395 And with its spectral type of B7/8 is particularly interesting because of its orbital period of only 0.685 days. FP Boo, SW Lac, V592 Per and TU UMi are contact binaries. Two of them, V592 Per and TU UMi, have bright tertiary companions, while FP Boo is a very small mass-ratio

($q = 0.11$) contact binary. The frequently observed system SW Lac is more massive than estimated before; its broadening functions show the large surface brightness and rather compact dimensions of its secondary component, both possibly indicating that it is really a semi-detached binary and not a contact system. Finally, we have been able to detect the low-mass secondary in FO Vir ($q = 0.12$); the system is most probably a semi-detached binary before establishing contact.

We express our thanks to Greg Stachowski for observations and Staszek Zola for advice on handling the case of KS Peg.

Special thanks go to Prof. Janusz Kaluzny who, acting as a referee, made very useful suggestions and comments which improved presentation of this paper.

Support from the Natural Sciences and Engineering Council of Canada to SMR and SWM and from the Polish Science Committee (KBN grants PO3D 006 22 and P03D 003 24) to WO and RP is acknowledged with gratitude.

The data given in this paper were obtained and analyzed when WP held the NATO Post-doctoral Fellowship administered by the Natural Sciences and Engineering Council of Canada.

The research made use of the SIMBAD database, operated at the CDS, Strasbourg, France and accessible through the Canadian Astronomy Data Centre, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada. This research made also use of the Washington Double Star (WDS) Catalog maintained at the U.S. Naval Observatory.

REFERENCES

- Abt, H. A. & Morrell, N. I. 1995, *ApJS*, 99, 135
- Adelman, S. J. 1998, *A&AS*, 132, 93
- Albayrak, B., Djurasevic, G., Erkapic, S. & Tanriverdi, T. 2004, *A&A*, 420, 1039
- Bessell, M. S. 1979, *PASP*, 91, 589
- Clausen, J. V., Helt, B. E. & Olsen, E. H. 2001, *A&A*, 374, 980
- Duerbeck, H. W. 1997, *Inf. Bull. Var. Stars*, 4513
- Duflot, M., Figon, P. & Meyssonier, N. 1995; *A&AS*, 114, 269
- Grenier S., Baylac M.O., Rolland L., Burnage R., Arenou F., Briot D., Delmas F., Duflot M., Genty V., Gomez A.E., Halbwachs J.-L., Marouard M., Oblak E., Sellier A. 1999, *A&AS*, 137, 451

- European Space Agency. 1997. The Hipparcos and Tycho Catalogues (ESA SP-1200)(Noordwijk: ESA) (HIP)
- Heintz, W. D. 1990, ApJS, 74, 275
- Hendry P. D. & Mochnacki, S. W. 1998, ApJ, 504, 978
- Høg, E., Fabricius, C., Makarov, V. V., Urban, S., Corbin, T., Wycoff, G., Bastian, U., Schwekendiek, P., & Wicenec, A. 2000, A&A, 355L, 27
- Hrivnak, B. J. 1992, Bull. AAS, 24, 686
- Hube, D. P. & Gulliver, A. F. 1985, PASP, 97, 280
- Hube, D. P., Martin, B. E. & Gulliver, A. F. 1988, Inf. Bull. Var. Stars, 3151
- Jordi, C., Figueras, F., Torra, J. & Asiain, R. 1996, A&AS, 115, 401
- Kazarovets, A. V., Samus, N. N., Durlevich, O. V., Frolov, M. S., Antipin, S. V., Kireeva, N. N., Pastukhova, E. N. 1999, Inf. Bull. Var. Stars, 4659
- Kreiner, J. M. 2004, Acta Astr., 54, 207
- Lu, W.-X., & Rucinski, S. M. 1999, AJ, 118, 515
- Lu, W., & Rucinski, S. M. 1993, AJ, 106, 361 (Paper I)
- Lu, W., Rucinski, S. M., & Ogloza, W. 2001, AJ, 122, 402 (Paper IV)
- Mason, B. D., Wycoff, G. L., Hartkopf, W. I., Douglass, G. G., & Worley, C. E. 2001, AJ, 122, 3466 (WDS)
- Mochnacki, S. W., Fernie, J. D., Lyons, R., Schmidt, F. H. & Gray, R. O. 1986, AJ, 91, 1221
- Morris, S. L. 1985, ApJ, 295, 143
- Pauzen, E. & Maitzen, H. M. 1998, A&AS, 133, 1
- Plaskett, J. S., Harper, W. E., Young, R. K., Plaskett, H. H. 1920, Publ. Dominion Astroph. Obs., 1, 163
- Popper, D.M. 1996, ApJS, 106,133
- Popper, D.M. 1997, AJ, 114, 1195
- Popper, D.M. 2000, AJ, 119, 2391
- Poretti, E., Niarchos, P. G., Mantegazza, L., Antonello, E. & Conconi, P. 1987, A&AS, 69, 337
- Pribulla, T., Chochol D., Parimucha, Š. 1999, Contr. Astron. Obs. Skalnaté Pleso, 29, 111

- Pribulla, T., Kreiner, J. M. & Tremko, J. 2003, *Contr. Astron. Obs. Skalnaté Pleso*, 33, 38
- Pych, W. & Rucinski, S. M. 2004, *Inf. Bull. Var. Stars*, 5524
- Pych, W., Rucinski, S. M., DeBond, H., Thomson, J. R., Capobianco, D. C., Blake, R. M., Ogloza, W., Stachowski, G., Rogoziecki, P., Ligeza, P., Gazeas, K. 2004, *AJ*, 127, 1712 (Paper IX)
- Rucinski, S. M. 2002a, *AJ*, 124, 1746 (Paper VII)
- Rucinski, S. M. 2002b, *PASP*, 114, 1124
- Rucinski, S. M., & Duerbeck, H.W. 1997, *PASP*, 109, 1340
- Rucinski, S. M., & Lu, W. 1999, *AJ*, 118, 2451 (Paper II)
- Rucinski, S. M., Lu, W., & Mochnacki, S. W. 2000, *AJ*, 120, 1133 (Paper III)
- Rucinski, S. M., Lu, W., Mochnacki, S. W., Ogloza, W., & Stachowski, G. 2001, *AJ*, 122, 1974 (Paper V)
- Rucinski, S. M., Lu, W., Capobianco, C. C., Mochnacki, S. W., Blake, R. M., Thomson, J. R., Ogloza, W., & Stachowski, G. 2002, *AJ*, 124, 1738 (Paper VI)
- Rucinski, S. M., Capobianco, C. C., Lu, W., DeBond, H., Thomson, J. R., Mochnacki, S. W., Blake, R. M., Ogloza, W., Stachowski, G., & Rogoziecki, P. 2003, *AJ*, 125, 3258 (Paper VIII)
- Schmidt, F. H. & Fernie, J. D. 1984, *Inf. Bull. Var. Stars*, 2527
- Selam, S. O. 2004, *A&A*, 416, 1097
- Shaw, J. S., Guinan, E. F. & Ivester, A. H. 1991, *Bull.AAS*, 23, 1415
- Zhai, D. & Lu, W. 1989, *Chinese Astr. Astroph.*, 13, 350

Captions to figures:

Fig. 1.— The broadening function “image” for V395 And with the orbital solution superimposed on the contour plot. The image has been smoothed in the phase domain with a Gaussian of FWHM = 0.025 in phase. A vertical section of the image at phase 0.75 is shown in Figure 5.

Fig. 2.— Radial velocities of the systems HS Aqr, V449 Aur, FP Boo and SW Lac are plotted in individual panels versus the orbital phases. The lines give the respective circular-orbit (sine-curve) fits to the radial velocities. HS Aqr and V449 Aur are detached binaries, while FP Boo and SW Lac are contact binaries. The circles and triangles in this and the next two figures correspond to components with velocities V_1 and V_2 , as listed in Table 1, respectively. The component eclipsed at the minimum corresponding to T_0 (as given in Table 2) is the one which shows negative velocities for the phase interval 0.0 – 0.5. The open symbols indicate observations contributing half (or less) weight in the orbital solutions. Short marks in the lower parts of the panels show phases of available observations which were not used in the solutions because of the blending of lines. All panels have the same vertical range, -350 to $+350$ km s $^{-1}$.

Fig. 3.— Same as in Figure 2, but for KS Peg, IW Per, V592 Per and FO Vir. Note that the vertical scale is different and much expanded for the single-line binaries KS Peg and IW Per, while for FO Vir it is the same for the other systems, but shifted vertically by $+50$ km s $^{-1}$.

Fig. 4.— The broadening function “image” for TU UMi with a very approximate guess at an orbital solution superimposed on the contour plot. The image has been smoothed in the phase domain with a Gaussian of FWHM = 0.025 in phase. Sections at phases 0.25 and 0.75 are shown in Figure 5.

Fig. 5.— The broadening function for all ten systems of this group. For most systems broadening functions at phases close to 0.25 have been selected. The phases are marked by numbers in individual panels; for KS Peg, FO Vir and TU UMi the numbers are written above locations of the secondary star peaks. For V395 And, KS Peg and TU UMi, the plots give actually sections of the “BF images”, after Gaussian smoothing (FWHM = 0.025) in the phase domain. For V395 And the section at phase 0.25 was too poorly defined so the section at phase 0.75 is shown instead. Note the high brightness of the secondary in SW Lac which directly shows the W-type phenomenon; note that this could be also interpreted by the semi-detached configuration of this binary (which otherwise is considered as one of the most typical contact systems). The uneven baseline for IW Per is due to differences in continuum tracing for the star and the template. As described in the text, FO Vir showed spurious peaks (one is visible here), which were almost as strong as those for the secondary component, but followed the motion of the primary component; we think it is because of the mismatch of the template with the low metallicity spectrum of FO Vir and because of the continuum height different for the template and for the star itself.

Table 1. DDO radial velocity observations (the full table is available only in electronic form)

HJD-2,400,000	V_1	W_1	V_2	W_2
HS Aqr				
2452507.6792	122.09	1.00	-143.05	1.00
2452507.7194	91.43	1.00	-104.07	1.00
2452507.7372	73.05	1.00	-99.45	0.50
2452507.7543	60.43	0.50	0.00	0.00
2452507.7744	38.30	0.50	0.00	0.00
2452507.7875	0.00	0.00	0.00	0.00
2452509.6398	84.74	1.00	-103.00	0.50
2452509.6551	95.57	1.00	-116.69	1.00
2452509.6704	108.27	1.00	-120.46	1.00
2452509.6863	115.38	1.00	-133.96	1.00

Note. — The table gives the radial velocities V_i and associated weights W_i for observations described in the paper. The first 10 rows of the table for the second program star, HS Aqr, are shown; this star is more typical than the first, V395 And, for which radial velocities were obtained from smoothed, averaged broadening functions as described in the text. There are no radial velocity data for TU UMi. In the table, velocities are expressed in km s^{-1} . Observations leading to entirely inseparable broadening- and correlation-function peaks are given zero weight; these observations may be eventually used in more extensive modeling of broadening functions. The radial velocities designated as V_1 correspond to the component which was stronger and easier to measure in the analysis of the broadening functions; it was not always the component eclipsed during the primary minimum at the epoch T_0 (see Table 2). The figures should help in identifying which star is which.

Table 2. Spectroscopic orbital elements

Name	Type Sp. type	Other names	V_0	K_1 K_2	ϵ_1 ϵ_2	$T_0 - 2,400,000$ (O–C)(d) [E]	P (days) ($M_1 + M_2$) $\sin^3 i$	q
V395 And ^a	EA B7/8:	HD 222900 HIP 117111	+14.60(1.51)	133.35(2.55) 151.72(2.06)	14.20 9.09	53,278.0000(30) +0.0332 [6978]	0.684656 1.647(80)	0.879(25)
HS Aqr	EA F6V	HD 197010 BD–01°4025	+19.34(0.64)	111.57(0.56) 178.24(1.46)	3.36 9.94	52,507.7981(12) +0.0002 [10]	0.710188 1.795(38)	0.626(7)
V449 Aur	EW(A) A2/3V	HD 41578 HIP 29108	+6.24(1.22)	128.27(1.45) 175.61(2.15)	9.72 16.76	52,307.0853(24) +0.0826 [5410]	0.703648 2.051(73)	0.730(15)
FP Boo	EW(A) F0V	BD+43°2523 HIP 76970	–4.87(1.02)	26.88(0.98) 254.04(2.80)	8.08 14.28	52,388.2227(36) –0.0114 [6070]	0.6404870 1.475(60)	0.106(5)
SW Lac	EW(W) G5V	HD 216598 HIP 113052	–10.32(1.17)	173.91(1.78) 224.14(1.70)	10.11 11.54	52,484.1080(8) +0.0003 [-50]	0.3207158 2.101(55)	0.776(12)
KS Peg	SB1 (EB?) A1V	HD 222133 HIP 116611	–3.27(0.62)	25.19(0.80)	5.25	50,735.6405(28) –0.0059 [16,115]	0.502103	
IW Per	SB1 (EB?) (A3V)	HD 21912 HIP 16591	+11.44(0.31)	98.64(0.51)	3.46	53,031.4192(8) +0.0270 4940]	0.91718	
V592 Per	EW(A) (F2IV)	HD 29911 HIP 22050	+27.92(1.04)	93.74(1.25) 229.97(1.85)	9.42 15.00	52,516.8641(20) –0.0108 [5612]	0.715722 2.521(72)	0.408(7)
TU UMi ^b	EW(W) (F2)	BD+76°544 HIP 73047	–4.16(0.20)	35(15) 220(20)	30 40	52,725.6262 0 [0]	0.377088 0.65(27)	0.16(7)
FO Vir	EA/EB (A7V)	HD 117362 HIP 65839	–27.26(1.45)	33.84(1.02) 269.91(2.95)	6.75 15.87	51,660.1839(25) –0.0342 [4074]	0.775568 2.257(89)	0.125(5)

^aV395 And: The solution is based on 103 data points arranged in phase using the preliminary T_0 and period, as given in the text, and then with the broadening functions smoothed in the phase domain. Such averaged data points are given in Table 1.

^bTU UMi: The data do not permit a solution; only very rough estimates are given. The value of V_0 is for the visual companion, while T_0 is from the photometric solution based on data obtained simultaneously with the spectroscopic program (Pych & Rucinski 2004). Because of the low contribution of the close binary to the total light, the (literature) spectral type is that of the visual companion.

Note. — The spectral types given in column two relate to the combined spectral type of all components in a system; they are given in parentheses if taken from the literature, otherwise are new. The convention of naming the binary components in the table is that the more massive star is marked by the subscript “1”, so that the mass ratio is defined to be always $q \leq 1$. The figures should help identify which component is eclipsed at the primary minimum. The standard errors of the circular solutions in the table are expressed in units of last decimal places quoted; they are given in parentheses after each value. The center-of-mass velocities (V_0), the velocity amplitudes (K_i) and the standard unit-weight errors of the solutions (ϵ) are all expressed in km s^{-1} . The spectroscopically determined moments of primary minima are given by T_0 ; the corresponding ($O - C$) deviations (in days) have been calculated from the available prediction on T_0 , as given in the text, using the assumed periods and the number of epochs given by [E]. The values of $(M_1 + M_2) \sin^3 i$ are in the solar mass units.









